

## PhD Thesis: development, validation and application of the TIM systems

- Arkbåge, K. (2003). Vitamin B12, folate and folate binding proteins in dairy products. Analysis, process retention and bioavailability. PhD Thesis Swedish University of Agricultural Sciences. Uppsala, Sweden. SLV Service/Repro. (ISBN: 91-567-6470-6).
- Krul, C.A.M. (2001, November 8). Mutagenic and antimutagenic activity of food compounds: application of a dynamic *in vitro* gastrointestinal model. PhD Thesis Utrecht University. Febodruk BV, Enschede, The Netherlands. (ISBN: 90-393-2857-9)
- Mateo Anson, N. (2010, May 28). Bioactive compounds in whole grain wheat. PhD Thesis, Maastricht University. Riddeprint b.v. The Netherlands. (ISBN: 978-90-5335-275-5).
- Minekus, M. (1998, May 28). Development and validation of a dynamic model of the gastrointestinal tract. PhD Thesis, Utrecht University. Elinkwijk b.v., Utrecht, Netherlands. (ISBN: 90-393-1666-x)
- Smeets-Peters, M.J.E. (2000, September 8). Feeding FIDO: Development, validation and application of a dynamic *in vitro* model of the gastrointestinal tract of the dog. PhD Thesis, Wageningen University. Universal Press, Veenendaal, The Netherlands. (ISBN: 90-5808-259-8).
- Van Nuenen, M. (2005, November 18). Composition and activity of gut microbiota in inflammatory bowel disease. PhD Thesis, Erasmus University Rotterdam. Ponsen & Looijen b.v., Wageningen, The Netherlands. (ISBN: 90-6464-234-6).
- Verwei, M. (2004, September 15). Bioavailability of folate from fortified milk products. PhD Thesis Wageningen University. The Netherlands (ISBN: 90-8504-080-9).

## General publications and technical aspects (incl. reviews)

- Abrahamse, E., Minekus, M., van Aken, G.A., van de Heijning, B., Knol, J., Bartke, N., Oozeer, R., van der Beek, E.M. and Ludwig, T. (2012). Development of the digestive system - Experimental challenges and approaches of infant lipid digestion. *Food Digestion* 3 (1-3): 63-77.
- Alminger, M., Aura, A.-M., Bohn, T., Dufour, C., El, S.N., Gomes, A., Karakaya, S., Martinez-Cuesta, M.C., McDougall, G.J., Requena, T., Santos, C.N. (2014). *In vitro* models for studying secondary plant metabolite digestion and bioaccessibility. *Compreh. Rev. Food Sci. Food Safety* 13: 413-436.
- Bellmann, S., Lelieveld, J., Gorissen, T., Minekus, M., Havenaar, R. (2016). Development of an advanced *in vitro* model and its evaluation versus human gastric physiology. *Food Res. Internat.* 88: 191-198.
- Culen, M., Rezacova, A., Jampilek, J., Dohnal, J. (2013). Designing a dynamic dissolution method: A review of instrumental options and corresponding physiology of stomach and small Intestine. *J. Pharmaceutical Sci.* DOI 10.1002/jps.23494.
- De Jong, L. (2005). Assessing Bioavailability. A new integrated methodology that predict long-term effects of foods in only a short time. *World Food Ingredients* Oct./Nov. pp 110-112.
- Freidig, A. and Verwei, M. (2004). Integration of *in vitro* data in kinetic models for pharmaceuticals and nutrients. *Netherlands Centre Altern. Animal Use Newsletter* 16: 1-3.
- Guerra, A., Etienne-Mesmin, L., Livrelli, V., Denis, S., Blanquet-Diot, S., Alric, M. (2012). Relevance and challenges in modeling human gastric and small intestinal digestion. *Trends Biotechnol.* 30 (11): 591-600.
- Havenaar, R., Veenstra, J., Minekus, M., Marteau, P. (1993). Unieke methode voor bestudering fysiologische aspecten van voeding. *Voeding* 54 (6): 7-11.
- Havenaar, R. and Minekus, M. (1996). Simulated assimilation. *Dairy Industries International* 61 (9): 17-23.
- Kostewicz ES, Abrahamsson B, Brewster M, Brouwers, J., Butler, J., Carlert, S., Dickinson, P.A., Dressman, J., Holm, R., Klein, S., Mann, J., McAllister, M., Minekus, M., Muenster, U., Müllertz, A., Verwei, M., Vertzoni, M., Weitschies, W., Augustijns, P. (2014). *In vitro* models for the prediction of *in vivo* performance of oral dosage forms. *Eur. J. Pharm. Sci.* 57 (1): 324-366.
- Li, Z., He, X. (2015). Physiologically based *in vitro* models to predict the oral dissolution and absorption of a solid drug delivery system. *Curr. Drug Metabol.* 16: 777-806.

- Lvova, L., Denis, S., Barra, A., Mielle, P., Salles, C., Vergoignan, C., Di Natale, C., Paolesse, R., Temple-Boyer, P. and Feron, G. (2012). Salt release monitoring with specific sensors in 'in vitro' oral and digestive environments from soft cheeses. *Talanta* 97: 171-180.
- Minekus, M. and Havenaar, R. (1998). Reactor system. European Patent No. 0642382. European Patent Bulletin 98/07, Art. 97(4) and (5) EPC, dated 11.02.98.
- Minekus, M. and Havenaar, R. (1996). In vitro model of an *in vivo* digestive tract. United States Patent; nr. 5,525,305, dated June 11, 1996.
- Minekus, M., Marteau, P., Havenaar, R., Huis in 't Veld, J.H.J. (1995). A multi compartmental dynamic computer-controlled model simulating the stomach and small intestine. *Alternatives to Laboratory Animals (ATLA)* 23: 197-209.
- Minekus, M., Smeets-Peeters, M.J.E., Bernalier, A., Marol-Bonnin, S., Havenaar, R., Marteau, P., Alric, M., Fonty, G., and Huis in 't Veld, J.H.J. (1999). A computer-controlled system to simulate conditions of the large intestine with peristaltic mixing, water absorption and absorption of fermentation products. *Appl. Microb. Biotechn.* 53: 108-114.
- Nguyen, T.T.P., Bhandari, B., Cichero, J., Prakash, S. (2015). A comprehensive review on *in vitro* digestion of infant formula. *Food Res. Internat.*
- Smeets-Peeters, M.J.E., Watson, T., Minekus, M., Havenaar, R. (1998). A review of the physiology of the canine digestive tract related to the development of *in vitro* systems. *Nutrition Research Reviews* 11: 45-69.
- Smeets-Peeters, M.J.E., Minekus, M., Havenaar, R., Schaafsma, G., Versteegen, M.W.A. (1999). Description of a dynamic *in vitro* model of the dog gastrointestinal tract and an evaluation of various transit times for protein and calcium. *ATLA* 27: 935-949.
- Ting, Y., Zhao, Q., Xia, C., Huang, Q. (2015). Using in vitro and in vivo models to evaluate the oral bioavailability of nutraceuticals. *J. Agr. Food Chem.* 63: 1332-1338.
- Venema, K. (2014). *In vitro* assessment of the bioactivity of food oligosaccharides. In: *Food Oligosaccharides: Production, Analysis and Bioactivity*. Eds.: F.J. Moreno & M. Luz Sanz. John Wiley & Sons, Ltd.
- Venema, K., Havenaar, R., Minekus, M. (2009). Improving *in vitro* simulation of the stomach and intestines. In: *Designing Functional Foods and Controlling Food Structure Breakdown and Nutrition Absorption*, Elsevier. pp 314-339.
- Verhoeckx, K. (2015). The impact of food bioactives on health: in vitro and ex-vivo models. Ed. Verhoeckx, K., Cotter, P., Lopez-Exposito, I. et al. Springer Open.
- Westerhout, J., van de Steeg, E., Grossouw, D., Zeijdner, E.E., Krul, C.A.M., Verwei, M., Wortelboer, H.M. (2014). A new approach to predict human intestinal absorption using porcine intestinal tissue and biorelevant matrices. *Eur. J. Pharmac. Sci.* 63: 167-177.
- Williams, C.F., Walton, G.E., Jiang, L., Plummer, S., Garaiova, I., Gibson, G.R. (2015). Comparative analysis of intestinal tract models. *Ann. Rev. Food Sci. Technol.* 6: 329-350.
- Yoo, J.Y., Chen, X.D. (2006). GIT Physicochemical modeling - A critical review. *Internat. J. Food Engineering*, 2 (4), article 4.
- Zhu Y. (2011) Response to the article "In vitro human digestion models for food application". *Food Chemistry*, 128: 820-821.

## Nutritional studies (human nutrition)

### Food Digestion (general)

- Keller, D., Van Dinter, R., Cash, H., Farmer, S., Venema, K. (2017). *Bacillus coagulans* GBI-30, 6086 increases plant protein digestion in a dynamic, computer-controlled *in vitro* model of the small intestine (TIM-1). *Beneficial Microbes* 8 (3): 491-496.
- Lefebvre, D.E., Venema, K., Gombau, L., Valerio Jr, L.G., Raju, J., Genevieve S. Bondy, G.S., Bouwmeester, H., Singh, R.P., Clippinger, A.J., Collnot, E.-M., Mehta, R., Stone, V. (2014). Utility of models of the gastrointestinal tract for assessment of the digestion and absorption of engineered

nanomaterials released from food matrices. *Nanotoxicology* (early on line 1-20; DOI: 10.3109/17435390.2014.948091).

Schnorr, .S.L., Crittenden, A.N., Venema, K., Marlowe, F., Henry, A.G. 2015. Assessing digestibility of Hadza tubers using a dynamic in-vitro model. *American Journal of Physical Anthropology* 158(3): 371-85.

## Proteins

Benmoussa, A., Lee, C.C., Laffont, B., Savard, P., Laugier, J., Boillard, E., Gilbert, C., Fliss, I., Provost, P. (2016) Commercial dairy cow milk microRNAs resist digestion under simulated gastrointestinal tract conditions. *J Nutr.* 146 (11): 2206-2215.

Denis, S., Sayd, T., Georges, A., Chambon, C., Chalancon, S., Santé-Lhoutellier, V., Blanquet-Diot, S. (2016). Digestion of cooked meat proteins is slightly affected by age as assessed using the dynamic gastrointestinal TIM model and mass spectrometry. *Food Function* 7 (6): 2682-2691.

Framroze, B., Savard, P., Gagnon, D., Richard, V., Gauthier, S.F. (2014). Comparison of nitrogen bioaccessibility from salmon and whey protein hydrolysates using a human gastrointestinal model (TIM-1). *Func. Foods Health Dis.* 4 (5): 222-231.

Havenaar, R., Maathuis, A., de Jong, A., Mancinelli, D., Berger, A., Bellmann, S. (2016). Herring roe protein had a high digestible indispensable amino acid score (DIAAS) using a dynamic *in vitro* gastrointestinal model. *Nutr. Res.* 36: 798-807.

Havenaar, R., de Jong, A., Koenen, M.J., van Bilsen, J., Janssen, A.M., Labij, E., Westerbeek, H.J.M. (2013). Digestibility of transglutaminase cross-linked caseinate versus native caseinate in an in vitro multi-compartmental model simulating young child and adult gastrointestinal conditions. *J. Agric. Food Chem.* 61 (31): 7636-7644.

Keller, D., Van Dinter, R., Cash, H., Farmer, S., Venema, K. (2017). *Bacillus coagulans* GBI-30, 6086 increases plant protein digestion in a dynamic, computer-controlled *in vitro* model of the small intestine (TIM-1). *Beneficial Microbes* 8 (3): 491-496.

Maathuis, A., Havenaar, R., He, T., Bellmann, S. (2017). Protein digestion and quality of goat and cow milk infant formula and human milk under simulated infant conditions. *J. Pediatric Gastroenterol. Nutr. Open Access.* doi:10.1097/MPG.0000000000001740.

Mitea, C., Havenaar, R., Drijfhout, J.W., Edens, L., Dekking, L. and Koning, F. (2008). Efficient degradation of gluten by prolyl endoprotease in gastrointestinal model: Implications for coeliac disease. *Gut* 57: 25-32.

Nabil, S., Gauthier, S.F., Drouin, R., Poubelle, P.E., and Pouliot, Y. (2011). In vitro digestion of proteins and growth factors in a bovine whey protein extract as determined using a computer-controlled dynamic gastrointestinal system (TIM-1). *Food Dig.* 2: 13-22.

Schaafsma, G. (2005). The Protein Digestibility-Corrected Amino Acid Score (PDCAAS). A concept for describing protein quality in foods and food ingredients: A critical review. *J. AOAC Internat.* 88 (3): 988-994.

Villemejeane, C., Denis, S., Marsset-Baglieri, A., Alric, M., Aymard, P., Michon, C. (2016). *In vitro* digestion of short-dough biscuits enriched in proteins and/or fibres, using a multi-compartmental and dynamic system (2): Protein and starch hydrolyses. *Food Chem.* 190: 164-172.

## Fats

Bellmann, S. und Havenaar, R. (2007). Die Fettbindenden Eigenschaften von Ballaststoffen *in vivo* und *in vitro*. *Ernährungs Umschau* 8/07: 450-455.

Bel-Rhliid, R., Pagé-Zoerkler, N., Fumeaux, R., Ho-Dac, T., Chuat, J-Y., Sauvageat, J.L., Raab, T. (2012). Hydrolysis of chicoric and caftaric acids with esterases and *Lactobacillus johnsonii* in vitro and in a gastrointestinal model. *J. Agric. Food Chem.* 60: 9236-9241.

Bel-Rhliid, R., Crespy, V., Pagé-Zoerkler, N., Nagy, K., Raab, T. and Hansen, C-E. (2009). Hydrolysis of rosmarinic acid from Rosemary extract with esterases and *Lactobacillus johnsonii* in vitro and in a gastrointestinal model. *J. Agric. Food Chem.* 57: 7700-7705.

- Domoto, N., Koenen, M.E., Havenaar, R., Mikajiri, A., Chu, B.-S., (2013). The bioaccessibility of eicosapentaenoic acid was higher from phospholipid food products than from mono- and triacylglycerol food products in a dynamic gastrointestinal model. *Food Sci. Nutr.* 1 (6): 409-415.
- Frank, N., Dubois, M., Scholz, G., Seefelder, W., Chuat, J.-Y., Schilter, B. (2013). Application of gastrointestinal modelling to the study of the digestion and transformation of dietary glycidyl esters. *Food Add. Containm. Part A*, 30 (1): 69-79. Doi: 10.1080/19440049.2012.732245.
- Fondaco, D., AlHasawi, F., Lan, Y., Ben-Elazar, S., Connolly, K., Rogers, M.A. (2015). Biophysical aspects of lipid digestion in human breast milk and Similac infant formulas. *Food Biophysics* 10: 282-291.
- Gervais, R., Gagnon, F., Kheadr, E.E., Van Calsteren, R.-M., Farnworth, E.R., Fliss, I., and Chouinard, P.Y. (2009). Bioaccessibility of fatty acids from conjugated linoleic acid-enriched milk and milk emulsions studied in a dynamic *in vitro* gastrointestinal model. *Internat. Dairy J.* 19: 574-581.
- Helbig, A., Silletti, E., van Aken, G.A., Oosterveld, A., Minekus, M., Hamer, R.J., Gruppen, H. (2013). Lipid digestion of protein stabilized emulsions investigated in a dynamic *in vitro* gastro-intestinal model system. *Food Dig.* 4: 58-68. (DOI 10.1007/s13228-012-0029-6).
- Larsson, K., Harrysson, H., Havenaar, R., Alminger, M., Undeland, I. (2016). Formation of malondialdehyde (MDA), 4-hydroxy-2-hexenal (HHE) and 4-hydroxy-2-nonenal (HNE) in fish and fish oil during dynamic gastrointestinal *in vitro* digestion. *Food Function* 7: 1176-1187. .
- Larsson, K., Tullberg, C., Alminger, M., Havenaar, R., Undeland, I. (2016). Malondialdehyde and 4-hydroxy-2-hexenal are formed during dynamic gastrointestinal *in vitro* digestion of cod liver oils. *Food Function* 7: 3458-3467.
- Maestre, R., Douglass, J.D., Kodukula, S., Medina, I, Storch, J. (2013). Alterations in the intestinal assimilation of oxidized PUFAs are ameliorated by a polyphenol-rich grape seed extract in an *in vitro* model and Caco-2 cells. *J. Nutr.* 143: 295-301.
- Minekus, M., Jelier, M., Xiao, J.-Z., Kondo S., Iwatsuki, K., Kokubo, S., Bos, M, Dunnewind, B. and Havenaar, R. (2005). Effect of partially hydrolyzed guar gum (PHGG) on the bioaccessibility of fat and cholesterol. *Biosci. Biotechnol. Biochem.* 69 (5): 932-938.
- Oosterveld, A., Minekus, M., Bomhof, E., Zoet, F.D., van Aken, G.A. (2016). Effects of inhomogeneity on triglyceride digestion of emulsions using an *in vitro* digestion model (tiny TIM). *Food Function* 7: 2979-2995.
- Reis, P.M., Raab, T.W., Chuant, J.Y., Leser, M., Miller, R., Watzke, H., and Holmberg, K. (2008). Influence of surfactants on lipase fat digestion in a model gastrointestinal system. *Food Biophysics* 3: 370-381.
- Speranza, A., Corradini, M.G., Hartman, T.G., Ribnicky, D., Oren, A., Rogers, M.A. (2013). Influence of emulsifier structure on lipid bioaccessibility in oil/water nanoemulsions. *J. Agr. Food Chem.* 61 (26): 6505-6515. (DOI: 10.1021/jf401548r).

### Carbohydrates (digestion and fermentation)

- Bellmann, S., Minekus, M., Sanders, P., Bosgra, S., Havenaar, R. (2017). Human glycemic response curves after intake of carbohydrate foods are accurately predicted by combining *in vitro* gastrointestinal digestion with *in silico* kinetic modeling. *Clin. Nutr. Experimental*. Open Access. doi.org/10.1016/j.yclnex.2017.10.003.
- Bothe M.K., Maathuis, A.J.H., Bellmann, S., van der Vossen, J.M.B.M., Berressem, D., Koehler, A., Schwejda-Guettes, S., Gaigg, B., Kuchinka-Koch, A., Stover, J.F. (2017). Dose-dependent prebiotic effect of lactulose in a computer-controlled *in vitro* model of the human large intestine. *Nutrients* 9: 767-781. (doi:10.3390.nu9070767).
- Faessler, C, Arrigoni, E., Venema, K., Hafner, V., Brouns, F. and Amado, R. (2006). Digestibility of resistant starch containing preparation using two *in vitro* models. *Eur. J. Nutr.* 45 (8): 445-453.
- Faessler, C, Arrigoni, E., Venema, K., Brouns, F. and Amado, R. (2006). *In vitro* fermentability of differently digested resistant starch preparations. *Mol. Nutr. Food Res.* 50: 1220-1228.
- Koenen, M.E., Cruz Rubio, J.M., Mueller, M., Venema, K. (2016). The effect of agave fructan products on the activity and composition of the microbiota determined in a dynamic *in vitro* model of the human proximal large intestine. *J. Funct. Foods* 22: 201-210.

- Lafond, M., Bouza, B., Eyrichine, S., Rouffineau, F., Saulnier, L., Giardina, T., Bonnin, E., Preynat, A. (2015). *In vitro* gastrointestinal digestion study of two wheat cultivars and evaluation of xylanase supplementation. *J. An. Sci. biotechnol.* 6 (1), Art. #5 (Open Access).
- Larsson Alminger, M., Eklund-Jonsson, C., Kidman, S., Langton, M. (2012). Starch microstructure and starch hydrolysis in barley and oat tempe during *in vitro* digestion. *Food Dig.* 3: 53-62.
- Maathuis, A.J.H., van den Heuvel, E.G., Schoterman, M.H.C., Venema, K. 2012. Galacto-Oligosaccharides have prebiotic activity in a dynamic *in vitro* colon model using a <sup>13</sup>C-labeling technique. *J. Nutrition* 142 (7): 1205-1212.
- Maathuis, A., Hoffman, A., Evans, A., Sanders, L., Venema, K. (2009). The Effect of the undigested fraction of maize products on the activity and composition of the microbiota determined in a dynamic *in vitro* model of the human proximal large intestine. *J. American College of Nutrition*, Vol. 28 (6): 657-666.
- Nalin, T. Venema, K., Weinstein, D.A., de Souza, C.F.M., Perry, I.D.S., van Wandelen, M.T.R., van Rijn, M., Smit, G.P.A., Schwartz, I.V.D., Derks, T.G.J. (2015). *In vitro* digestion of starches in a dynamic gastrointestinal model: an innovative study to optimize dietary management of patients with hepatic glycogen storage diseases. *J. Inherit. Metab. Dis.* 38: 529-536. Erratum: *J. Inherit. Metab. Dis.* 38 (5): 987.
- Ramasamy, U.S., Venema, K., Schols, H.A., Gruppen, H. (2014). Effect of soluble and insoluble fibers within the *in vitro* fermentation of chicory root pulp by human gut bacteria. *J. Agr. Food Chem.* 62: 6794-6802.
- Ramasamy, U.S., Venema, K., Gruppen, H., Schols, H.A. (2014). The fate of chicory root pulp polysaccharides during fermentation in the TNO *in vitro* model of the colon (TIM-2). *Bioact. Carb. Dietary Fibre* 4: 48-57.
- Reimer, R.A., Maathuis, A.J.H., Venema, K., Lyon, M.R., Gahler, R.J., Wood, S. (2014). Effect of the novel polysaccharide PolyGlycopleX® on short-chain fatty acid production in a computer-controlled *in vitro* model of the human large intestine. *Nutrients* 6: 1115-1127. (doi: 10-3390/nu6031115).
- Van Nuenen, H.M.C., Meyer, P.D., Venema, K. (2003). The effect of various inulins and *Clostridium difficile* on the metabolic activity of the human colonic microbiota *in vitro*. *Microbial Ecology in Health and Disease* 15 (2-3): 137-144.
- Venema, K., Van Nuenen, H.M.C., Van den Heuvel, E.G., Pool, W., Van der Vossen, J.M.B.M. (2003). The effect of lactulose on the composition of the intestinal microbiota and short-chain fatty acid production in human volunteers and a computer-controlled model of the proximal large intestine. *Microbial Ecology in Health and Disease*, 15 (2-3): 94-105.
- Villemejeane, C., Denis, S., Marsset-Baglieri, A., Alric, M., Aymard, P., Michon, C. (2016). *In vitro* digestion of short-dough biscuits enriched in proteins and/or fibres, using a multi-compartmental and dynamic system (2): Protein and starch hydrolyses. *Food Chem.* 190: 164-172.
- Villemejeane, C., Wahl, R., Aymard, P., Denis, S., Michon, C. (2015). *In vitro* digestion of short-dough biscuits enriched in proteins and/or fibres, using a multi-compartmental and dynamic system (1): Viscosity measurement and prediction. *Food Chem.* 182: 55-63.

## Vitamins

- Arkbåge, K., Verwei, M., Havenaar, R., Witthöft, C. (2003). Folic acid and (6S)-5-methyltetrahydrofolate bioaccessibility decreases after addition of folate-binding protein to yogurt as studied in a dynamic *in vitro* gastrointestinal model. *J. Nutr.* 133: 3678-3683.
- Blanquet-Diot, S., Soufi, M., Rambeau, M., Rock, E., and Alric, M. (2009). Digestive stability of xanthophylls exceeds that of carotenes as studied in a dynamic *in vitro* gastrointestinal system, *J. Nutr.* 139 (5): 876-883.
- Déat, E., Blanquet-Diot, S., Jarrige, J-F., Denis, S., Beyssac, E. and Alric, M. (2009). Combining the dynamic TNO-gastrointestinal tract system with a Caco-2 cell culture model: Application to the assessment of lycopene and  $\alpha$ -tocopherol bioavailability from a whole food. *J. Agricult. Food Chem.* 57: 11314-11320. (Correction of Fig. 4: *JAF* p 11314).

- Etcheverry, P., Grusak, M.A., Fleige, L.E. (2012). Application of *in vitro* bioaccessibility and bioavailability methods for calcium, carotenoids, folate, iron, magnesium, polyphenols, zinc, and vitamins B6, B12, D, and E. *Frontiers Physiology* p1-22; doi: 10.3389/fphys.2012.00317.
- Finglas, P.M., de Meer, K., Molloy, A., Verhoef, P., Pietrzik, K., Powers, H.J., van der Straeten, D., Jägerstad, M., Varela-Moreiras, G., van Vliet, T., Havenaar, R., Buttriss, J., Wright, A.J.A. (2007). Research goals for folate and related B vitamin in Europe. *Eur. J. Clin. Nutr.* 60 (2): 287-294.
- Galán, I., García, M.L., Selgas, M.D., Havenaar, R. (2014). Effect of E-beam treatment on the bioaccessibility of folic acid incorporated to ready to eat meat products. *Food Sci. Techn.* 59: 547-552.
- Ohrvik, V., Witthöft, C., (2008). Orange juice is a good folate source in respect to folate content and stability during storage and simulated digestion. *Eur. J. Nutr.* 47: 92-98.
- Ohrvik, V., Ohrvik, H., Tallkvist, J., Witthöft, C. (2010). Foliates in bread: retention during bread-making and *in vitro* bioaccessibility. *Eur. J. Nutr.* 49 (6): 365-372.
- Nimalaratne, C., Savard, P., Gauthier, S.F., Schieber, A., Wu, J. (2015). Bioaccessibility and digestive stability of carotenoids in cooked eggs studied using a dynamic *in vitro* gastrointestinal model. *J. Agr. Food Chem.* 63 (11): 2956-2962.
- Richelle, M., Sanchez, B., Tavazzi, I., Lambelet, P., Bortlik, K. and Williamson, G. (2010). Lycopene isomerisation takes place within enterocytes during absorption in human subjects. *Br. J. Nutr.* 103: 1800-1807.
- Van Loo-Bouwman, C.A., Naber, T.H.J., Minekus, M., van Breemen, R.B., Hulshof, P.J.M., Schaafsma, G. (2014). Food matrix effects on bioaccessibility of  $\beta$ -carotene can be measured in an *in vitro* gastrointestinal model. *J. Agric. Food Chem.* 62 (4): 950-955.
- Verwei, M., Freidig, A.P., Havenaar, R., Groten, J. P. (2006). Predicted serum folate concentrations based on *in vitro* studies and kinetic modeling are consistent with measured folate concentrations in humans. *J. Nutr.* 136 (12): 3074-3078.
- Verwei, M., Arkbåge, K., Groten, J.P., Witthöft, C., Van den Berg, H. and Havenaar, R. (2005). The effect of folate binding proteins on bioavailability of folate from milk products. *Trends Food Sci. Techn.* 16: 307-310.
- Verwei, M., Arkbåge, K., Mocking, H., Havenaar, R. and Groten, J. (2004). The binding of folic acid and 5-methyltetrahydrofolate to folate-binding proteins during gastric passage differs in a dynamic *in vitro* gastrointestinal model. *J. Nutr.* 134: 31-37.
- Verwei, M., Arkbåge, K., Havenaar, R., Van den Berg, H., Witthöft, C. and Schaafsma, G. (2003). Folic acid and 5-Methyl-tetrahydrofolate in fortified milk are bioaccessible as determined in a dynamic *in vitro* gastrointestinal model. *J. Nutr.* 133: 2377-2383.

## Minerals

- Bellmann, S., Miyazaki, K., Chonan, O., Ishikawa, F., Havenaar, R. (2014). Fucoidan from *Cladosiphon okamuranus* Tokida added to food has no adverse effect on availability for absorption of divalent minerals in the dynamic multi-compartmental model of the upper gastrointestinal tract. *Food Digestion* 5 (1-3): 19-25. DOI 10.1007/s13228-014-0036-x
- Eklund-Jonsson, C., Sandberg, A-S., Hulthen, L., Larsson-Alminger, M. (2008). Tempé fermentation of whole grain barley increased human iron absorption and *in vitro* iron availability. *The Open Nutr. J.* 2: 42-47.
- Etcheverry, P., Grusak, M.A., Fleige, L.E. (2012). Application of *in vitro* bioaccessibility and bioavailability methods for calcium, carotenoids, folate, iron, magnesium, polyphenols, zinc, and vitamins B6, B12, D, and E. *Frontiers Physiology* p1-22; doi: 10.3389/fphys.2012.00317.
- Haraldsson, A-K., Rimsten, L., Alminger, M., Andersson, R., Aman, P., and Sandberg, A-S. (2005). Digestion of barley malt porridges in a gastrointestinal model: Iron dialysability, iron uptake by Caco-2 cells and degradation of  $\beta$ -glucan. *J. Cereal Sci.* 42: 243-254.
- Kortman, G.A.M., Dutilh, B.E., Maathuis, A.J.H., Engelke, U.F., Boekhorst, J., Keegan, K.P., Nielsen, F.G.G., Betley, J., Weir, J.C., Kingsbury, Z., Kluijtmans, L.A.J., Swinkels, D.W., Venema, K., Tjalsma, H. (2016). Microbial metabolism shifts towards an adverse profile with supplementary iron

- in the TIM-2 *in vitro* model of the human colon. *Frontiers Microbiol.* 6, art. 1481. doi: 10.3389/fmicb.2015.01481.
- Larsson, M. Minekus, M. and Havenaar, R. (1997). Estimation of the bio-availability of iron and phosphorus in cereals using a dynamic *in vitro* gastrointestinal model. *J. Sci. Food Agric.* 73: 99-106.
- Lvova, L., Denis, S., Barra, A., Mielle, P., Salles, C., Vergoignan, C., Di Natale, C., Paolesse, R., Temple-Boyer, P. and Feron, G. (2012). Salt release monitoring with specific sensors in '*in vitro*' oral and digestive environments from soft cheeses. *Talanta* 97: 171-180.
- Martin, A.H. and De Jong, G.A.H. (2012). Enhancing the *in vitro* Fe<sup>2+</sup> bioaccessibility using ascorbate and cold-set whey protein gel particles. *Dairy Sci. Technology* 92 (2): 133-149.
- Salovaara, S., Larsson-Alminger, M., Eklund-Jonsson, C., Andlid, T. and Sandberg, A.-S. (2003). Prolonged transit time through the stomach and small intestine improves iron dialyzability and uptake *in vitro*. *J. Agric. Food Chem.* 51: 5131-5136.

## Antioxidants

- Hemery, Y.M., Mateo Anson, N., Havenaar, R., Haenen, G.R.M.M., Noort, M.W.J. Rouau, X. (2010). Dry-fractionation of wheat bran increases the bioaccessibility of phenolic acids in breads made from processed bran fractions. *Food Research International*, 43 (5): 1429-1438.
- Lafond, M., Bouza, B., Eyriche, S., Bonnin, B., Crost, E.H., Geraert, P-A., Giardina, T., and Ajandouz E.H. (2011). An integrative *in vitro* approach to analyse digestion of wheat polysaccharides and the effect of enzyme supplementation. *British J. Nutr.* 106: 264-273.
- Lila, M.A., Ribnicky, D.M., Rojo, L.E., Rojas-Silva, P., Oren, A., Havenaar, R., Janle, E.M., Raskin, I., Yousef, G.G., and Grace, M.H. (2012). Complementary approaches to gauge the bioavailability and distribution of ingested berry polyphenols. *J. Agricultural Food Chem.* 60: 5763-5771.
- Maestre, R., Douglass, J.D., Kodukula, S., Medina, I, Storch, J. (2013). Alterations in the intestinal assimilation of oxidized PUFAs are ameliorated by a polyphenol-rich grape seed extract in an *in vitro* model and Caco-2 cells. *J. Nutr.* 143: 295-301.
- Mateo Anson, N.; Havenaar, R.; Bast, A. and Haenen, G.R.M.M. (2010). Antioxidant and anti-inflammatory capacity of bioaccessible compounds from wheat fractions after gastrointestinal digestion. *J. Cereal Sci.* 51 (1): 110-114.
- Mateo Anson, N., Van den Berg, R., Havenaar, R., Bast, A., Haenen, G. (2009). Bioavailability of ferulic acid is determined by its bioaccessibility. *J. Cereal Sci.* 49 (2): 295-300.
- Mateo Anson, N, Selinheimo, E., Havenaar, R., Aura, A.M., Mattila, I., Lehtinen, P., Bast, A., Poutanen, K., Haenen, G.R.M.M. (2009). Bioprocessing of wheat bran improves *in vitro* bioaccessibility and colonic metabolism of phenolic compounds. *J. Agric. Food Chem.* 57: 6148-6155.
- Mateo Anson, N., Havenaar, R., Vaes, W., Coulier, L., Venema, K., Selinheimo, E., Bast, A., Haenen, G.R.M.M. (2011). Effect of bioprocessing of wheat bran in whole meal wheat breads on the colonic SCFA production *in vitro* and postprandial plasma concentrations in men. *Food Chemistry* 128: 404-409.

## Animal feed & nutrition

- Avantaggiato, G., Havenaar, R. and Visconti, A. (2007). Assessment of the multi-mycotoxin binding efficacy of a carbon/aluminosilicate based product in an *in vitro* gastrointestinal model. *J. Agricul. Food Chem.* 55: 4810-4819.
- Avantaggiato, G., Havenaar, R. and Visconti, A. (2004). Evaluation of the intestinal absorption of deoxynivalenol and nivalenol by an *in vitro* gastrointestinal model, and the binding efficacy of activated carbon and other adsorbent materials. *Food Chemical Tox.* 42 (5): 817-824.
- Avantaggiato, G., Havenaar, R. and Visconti, A. (2003). Assessing the zearalenone binding activity of adsorbent materials during passage through a dynamic gastrointestinal model. *Food Chemical Tox.* 41: 1283-1290.

- Havenaar, R. (2002). GI Model. FIDO: An alternative to invasive animal research. *Petfood Industry* 44: 12-18.
- Lankhorst, C., Tran, Q., Havenaar, R., Hendriks, W., van der Poel, A. (2007). The effect of extrusion on the nutritional value of canine diets as assessed by *in vitro* indicators. *Animal Feed Sci. Techn.* 138: 285-297.
- Meunier, J.-P., Cardot, J.M., Manzanilla, E.G., Wysshaar, M., and Alric, M. (2007). Use of spray-cooling technology for development of microencapsulated capsaicin oleoresin for the growing pig as an alternative to in-feed antibiotics: A study of release using *in vitro* models. *J. Anim. Sci.* 86: 1156-1163.
- Meunier, JP, Manzanilla, EG, Anguita, M, Denis, S, Perez, JF, Gasa, J, Cardot, J-M,, Garcia, F, Moll, X, Alric, M.. (2008). Evaluation of a dynamic *in vitro* model to simulate the porcine ileal digestion of diets differing in carbohydrate composition. *J. Anim Sci.* 86: 1156-1163.
- Minekus, M. (1996). *In-vitro* modelling moves closer to the real thing. *Feed Mix* 4: 36-38.
- Schaafsma, G. (2005). The Protein Digestibility-Corrected Amino Acid Score (PDCAAS). A concept for describing protein quality in foods and food ingredients: A critical review. *J. AOAC Internat.* 88 (3): 988-994.
- Smeets-Peters, M.J.E., Minekus, M., Havenaar, R., Schaafsma, G., Verstegen, M.W.A. (1999). Description of a dynamic *in vitro* model of the dog gastrointestinal tract and an evaluation of various transit times for protein and calcium. *ATLA* 27: 935-949.

## Other nutrition studies

- Björck, I., Ostman, E., Kristensen, M., Mateo Anson, N., Price, R., Haenen, G., Havenaar, R., Bach Knudsen, K-E., Frid, A., Mykkanen, H., Welch, R., and Riccardi, G. (2012). Cereal grains for nutrition and health benefits: Overview of results from *in vitro*, animal and human studies in the HEALTHGRAIN Project. *Trends in Food Science & Technology* 25 (2): 87-100.
- Helou, C., Denis, S., Spatz, M., Marier, D., Rame, V., Alric, M., Tessier, F.J., Gadonna-Widehem, P. (2015). Insights into bread melanoidins: Fate in the upper digestive tract and impact on the gut microbiota using *in vitro* systems. *Food Func.* 6 (12): 3737-3745.
- Nguyen, T.T.P., Bhandari, B., Cichero, J., Prakash, S. (2015). A comprehensive review on *in vitro* digestion of infant formula. *Food Res. Internat.*

## Pharmaceutical studies

- Barker, R., Abrahamsson, B., Kruusmägi, M. (2014). Application and validation of an advanced gastrointestinal *in vitro* model for evaluation of drug product performance in pharmaceutical development. *J. Pharm. Sci.* 103 (11): 3704-3712. DOI 10.1002/jps.24177.
- Blanquet, S.; Garrait, G.; Beyssac, E.; Perrier, C.; Denis, S.; Hebrard, G.; Alric, M. (2005). Effects of cryoprotectants on the viability and activity of freeze dried recombinant yeasts as novel oral drug delivery systems assessed by an artificial digestive system. *Eur. J. Pharmaceutics and Biopharmaceutics*, 61: 32-39.
- Blanquet, S., Antonelli, R., Laforet, L., Denis, S., Marol-Bonin, S. and Alric, M. (2004). Living recombinant *Saccharomyces cerevisiae* secreting proteins or peptides as a new drug delivery system in the gut. *J. Biotechnol.* 110: 37-49.
- Blanquet, S., Zeijdner, E., Beyssac, E., Meunier, J-P., Denis, S., Havenaar, R. and Alric, M. (2004). A dynamic artificial gastrointestinal system for studying the behavior of orally administered drug dosage forms under various physiological conditions. *Pharmaceutical Research*. 21 (4): 585-591.
- Blanquet, S., Marol-Bonin, S., Beyssac, E., Pompon, D., Renaud, M. and Alric, M. (2001). The 'biodrug' concept: an innovative approach to therapy. *Trends Biotechnol.* 19 (10): 393-400.
- Brouwers, J., Anneveld, B., Goudappel, G.J, Duchateau, G., Annaert, P., Augustijns, P. and Zeijdner, E. (2011) Food-dependent disintegration of immediate release fosamprenavir tablets: *In vitro*



- evaluation using magnetic resonance imaging and a dynamic gastrointestinal system. *Eur. J. Pharmaceutics Biopharmaceutics*, 77: 313–319.
- Chen, L., Hebrard, G., Beyssac, E., Denis, S. and Subirade, M. (2010). *In vitro* study of the release properties of soy-zein protein microspheres with a dynamic artificial digestive system. *J. Agricultural Food Chem.* 58: 9861–9867.
- David, S.E., Strozyk, M.M. & Naylor, T.A. (2010). Using TNO gastro-Intestinal Model (TIM-1) to screen potential formulations for a poorly soluble development compound. *J. Pharm. Pharmacol.* 62: 1236–1237.
- De Almeida, M.R.A., Bassani, A.S., Hamid, S., Banov, D., Phan, H. (2015). Investigation of the bioaccessibility of progesterone (micronized and non-micronized), using an *in vitro* model of the human gastrointestinal system. *Intern. J. Appl. Sci. Techn.* 5 (4): 1-7.
- Dickinson, P.A., Abu Rmaileh, R., Ashworth, L., Barker, R.A., Burke, W.M., Patterson, C.M., Stainforth, N. and Yasin, M. (2012). An investigation into the utility of a multi-compartmental, dynamic, system of the upper gastrointestinal tract to support formulation development and establish bioequivalence of poorly soluble drugs. *AAPS Journal* 14 (2): 196-205.
- Garbacz, G., Klein, S. (2012). Dissolution testing of oral modified-release dosage forms. *J. Pharmacy Pharmacology* 64: 944-958.
- Gittings, S., Turnbull, N., Roberts, C.J., Gershkovich, P. (2014). Dissolution methodology for taste masked oral dosage forms. *J. Controlled Release* 173: 32-42.
- Havenaar, R., Anneveld, B., Hanff, L.M., de Wildt, S.N., de Koning, B.A.E., Mooij, M.G., Lelieveld, J.P.A., Minekus, M. (2013). In vitro gastrointestinal model (TIM) with predictive power, even for infants and children? *Internat. J. Pharm.* 457: 327-332.
- Hens, B., Brouwers, J., Anneveld, B., Corsetti, M., Symillides, M., Vertzoni, M., Reppas, C., Turner, D.B., Augustijns, P. (2014). Gastrointestinal transfer: *In vivo* evaluation and implementation in *in vitro* and *in silico* predictive tools. *Eur. J. Pharmaceut. Sci.* 63: 233-242.
- Kostewicz ES, Abrahamsson B, Brewster M, Brouwers, J., Butler, J., Carlert, S., Dickinson, P.A., Dressman, J., Holm, R., Klein, S., Mann, J., McAllister, M., Minekus, M., Muenster, U., Müllertz, A., Verwei, M., Vertzoni, M., Weitschies, W., Augustijns, P. (2014). In vitro models for the prediction of in vivo performance of oral dosage forms. *Eur. J. Pharm. Sci.* 57 (1): 342-366.
- Koziolek, M., Garbacz, G., Neumann, M., Weitschies, W. (2013). Simulating the postprandial stomach: Biorelevant test methods for the estimation of intragastric drug dissolution. *Mol. Pharmaceutics* 10: 2211-2221.
- Lyng, E., Havenaar, R., Shastri, P., Hetsco, L., Vick, A., Sagartz, J. (2016). Increased bioavailability of celecoxib under fed versus fasted conditions is determined by post-prandial bile secretion as demonstrated in a dynamic gastrointestinal model. *J. Drug Dev. Ind. Pharm.* 42 b(8): 1334-1339.
- McAllister, M. (2010). Dynamic Dissolution: A Step Closer to Predictive Dissolution Testing? *Molecular Pharmaceutics* 7 (5): 1374-1387.
- Naylor, T.A., Connolly, P.C., Martini, L.G., Elder, D.P., Minekus, M., Havenaar, R. and Zeijdner, E. (2006). Use of a gastro-intestinal model and Gastroplus™ for the prediction of *in vivo* performance. *Industrial Pharmacy, Issue 12*: 9-12. (also published in: *Applied Therapeutic Research* 6 (1): 15-19.
- Rogers, M.A., Yan, Y.-F., Ben-Elazar, K., Lan, Y., Faig, J., Smith, K., Uhrich, K.E. (2014). Salicylic Acid (SA) Bioaccessibility from SA-Based Poly(anhydrideester). *Biomacromolecules* 15: 3406–3411.
- Salmon, F., Verwei, M., Havenaar, R. (2008). When oral bioavailability becomes a problem. *Eur. Biopharm. Rev.*: 78-83.
- Souliman, S., Beyssac, E., Cardot, J.-M., Denis, S. and Alric, M. (2007). Investigation of the biopharmaceutical behavior of theophylline hydrophilic matrix tablets using USP methods and an artificial digestive system. *Drug Development & Industrial Pharm.* 33 (4): 475-483.
- Souliman, S., Blanquet, S., Beysac, E. and Cardot, J.-M. (2006). A level A *in vitro/in vivo* correlation in fasted and fed states using different methods: Applied to solid immediate release oral dosage form. *Eur. J. Pharmaceutical Sci.* 27: 72-79.

- Tenjarla S, Romasanta V, Zeijdner E, Villa R, Moro L. (2007). Release of 5-aminosalicylate from an MMX mesalamine tablet during transit through a simulated gastrointestinal tract system. *Adv Ther.* 24 (4): 826-840.
- Ting, Y., Jiang, Y., Lan, Y., Lin, Z., Rogers, M.A., Huang, Q. (2015). Viscoelastic emulsion improved the bioaccessibility and oral bioavailability of crystalline compound: A mechanistic study using *in vitro* and *in vivo* models. *Mol. Pharmaceutics* 12 (7): 2229-2236.
- Van Den Abeele, J., Schilderink, R., Schneider, F., Mols, R., Minekus, M., Weitschies, W., Brouwers, J., Tack, J., Augustijns, P. (2017). Gastrointestinal and systemic disposition of diclofenac under fasted and fed state conditions supporting the evaluation of *in vitro* predictive tools. *Mol. Pharmaceutics* doi: 10.1021/acs.molpharmaceut.7b00253.
- Verwei, M., Minekus, M., Zeijdner, E., Schilderink, R., Havenaar, R. (2016). Evaluation of two dynamic *in vitro* models simulating fasted and fed state conditions in the upper gastrointestinal tract (TIM-1 and tiny-TIM) for investigating the bioaccessibility of pharmaceutical compounds from oral dosage forms. *Int. J. Pharm.* 498: 178-186.
- Westerhout, J., van de Steeg, E., Grossouw, D., Zeijdner, E.E., Krul, C.A.M., Verwei, M., Wortelboer, H.M. (2014). A new approach to predict human intestinal absorption using porcine intestinal tissue and biorelevant matrices. *Eur. J. Pharmac. Sci.* 63: 167-177.
- Zeijdner, E.E., Vlek, J. (2002). TIM: a versatile tool in studying paediatric pharmacokinetics. *The Regulatory Review (The Journal of the British Institute of Regulatory Affairs)* 5 (7):18-21.
- Zeijdner, E.E. and Havenaar, R. (2000). The fate of orally administered compounds during passage through the gastrointestinal tract simulated in a dynamic *in vitro* model (TIM). *European Pharmaceutical Contractor*, Febr. issue: 76-81.

## Functional Ingredients, Phytochemicals & Biologicals

- Benmoussa, A., Lee, C.C., Laffont, B., Savard, P., Laugier, J., Boilard, E., Gilbert, C., Fliss, I., Provost, P. (2016). Commercial dairy cow milk microRNAs resist digestion under simulated gastrointestinal tract conditions. *J Nutr.* 146 (11): 2206-2215.
- Bothe M.K., Maathuis, A.J.H., Bellmann, S., van der Vossen, J.M.B.M., Berressem, D., Koehler, A., Schwejda-Guettes, S., Gaigg, B., Kuchinka-Koch, A., Stover, J.F. (2017). Dose-dependent prebiotic effect of lactulose in a computer-controlled *in vitro* model of the human large intestine. *Nutrients* 9: 767-781. (doi:10.3390.nu9070767).
- Gagnon, M., Savard, P., Rivière, A., LaPointe, G., Roy, D. (2015). Bioaccessible antioxidants in milk fermented by *Bifidobacterium longum* subsp. *longum* strains. *BioMed Res. Internat.*, ID 169381. <http://dx.doi.org/10.1155/2015/169381>.
- Gao, K., Xu, A., Krul, C., Venema, K., Liu, Y., Niu, Y., Lu, J., Bensoussan, L., Seeram, N.P., Heber, D. and Henning, S.M. (2006). Of the major phenolic acids formed during human microbial fermentation of tea, citrus and soy flavonoids supplements, only 3,4-dihydroxyphenylacetic acid has antiproliferative activity. *J. Nutr.* 136: 52-57.
- Krul, C., Humblot, C., Philippe, C., Vermeulen, M., Van Nuenen, M., Havenaar, R., Rabot, S. (2002). Metabolism of sinigrin (2-propenyl glucosinolate) by the human colonic microflora in a dynamic *in vitro* large-intestinal model. *Carcinogenesis* 23 (6): 1009-1016.
- Krul, C.A.M., Luiten-Schuite, A., Tenfelde, A., Van Ommen, B., Verhagen, H. and Havenaar, R. (2001). Antimutagenic activity of green tea and black tea extracts studied in a dynamic *in vitro* gastrointestinal model. *Mutation Research* 474: 71-85.
- Kong, H., Wang, M., Venema, K., Maathuis, A., van der Heijden, R., van der Greef, J., Xu, G. and Hankemeier, T. (2009). Bioconversion of red ginseng saponins in the gastro-intestinal tract *in vitro* model studied by high-performance liquid chromatography-high resolution Fourier transform ion cyclotron resonance mass spectrometry. *J. Chromatography A* 1216: 2195-2203.
- Kheadr, E., Zihler, A., Dabour, N., Lacroix, C., Le Blay, G. and Fliss, I. (2010). Study of the physicochemical and biological stability of pediocin PA-1 in the upper gastrointestinal tract conditions using a dynamic *in vitro* model. *J. Appl. Microbiol.* 109: 54-64.

- Lila, M.A., Ribnicky, D.M., Rojo, L.E., Rojas-Silva, P., Oren, A., Havenaar, R., Janle, E.M., Raskin, I., Yousef, G.G., and Grace, M.H. (2012). Complementary approaches to gauge the bioavailability and distribution of ingested berry polyphenols. *J. Agricultural Food Chem.* 60: 5763-5771.
- Maestre, R., Douglass, J.D., Kodukula, S., Medina, I., Storch, J. (2013). Alterations in the intestinal assimilation of oxidized PUFAs are ameliorated by a polyphenol-rich grape seed extract in an in vitro model and Caco-2 cells. *J. Nutr.* 143: 295-301.
- Mitea, C., Havenaar, R., Drijfhout, J.W., Edens, L., Dekking, L. and Koning, F. (2008). Efficient degradation of gluten by prolyl endoprotease in a gastrointestinal model: Implications for coeliac disease. *Gut* 57: 25-32.
- Minekus, M., Jelier, M., Xiao, J.-Z., Kondo S., Iwatsuki, K., Kokubo, S., Bos, M, Dunnewind, B. and Havenaar, R. (2005). Effect of partially hydrolyzed guar gum (PHGG) on the bioaccessibility of fat and cholesterol. *Biosci. Biotechnol. Biochem.* 69 (5): 932-938.
- Nabil, S., Gauthier, S.F., Drouin, R., Poubelle, P.E., and Pouliot, Y. (2011). In vitro digestion of proteins and growth factors in a bovine whey protein extract as determined using a computer-controlled dynamic gastrointestinal system (TIM-1). *Food Dig.* 2: 13-22.
- Ribnicky, D.M., Roopchand, D.E., Poulev, A., Kuhn, P., Oren, A., Cefalu, W.T., Raskin, I. (2014). *Artemisia dracunculoides* L. polyphenols complexed to soy protein show enhanced bioavailability and hypoglycemic activity in C57BL/6 mice. *Nutrition* 30: S4-S10.
- Ribnicky, D.M., Roopchand, D.E., Oren, A., Grace, M., Poulev, A., Lila, M.A., Havenaar, R., Raskin, I. (2014). Effects of a high fat meal matrix and protein complexation 4 on the bioaccessibility of blueberry anthocyanins using the TNO gastrointestinal model (TIM-1). *Food Chem.* 142: 349-357.
- Thörig, L., de Groot, L., Hensgens, C.M.H. (2001). Bovine hyperimmune whey protein concentrate with specific biological activity as a replacement ingredient. *Innovations Food Techn.* Dec.2001: 57-60.

## Probiotics & Microbiology studies (see also micro-ecology TIM-2 studies)

- Arroyo-López, F.N., Blanquet-Diot, S, Denis, S., Thévenot, J., Chalancon, S., Alric, M., Rodríguez-Gómez, F., Romero-Gil, V., Jiménez-Díaz, R., Garrido-Fernández, A. (2014). Survival of pathogenic and lactobacilli species of fermented olives during simulated human digestion. *Frontiers Microbiol.* p1-9 doi 10.3389/micb.2014.00540.
- Bel-Rhliid, R., Pagé-Zoerkler, N., Fumeaux, R., Ho-Dac, T., Chuat, J-Y., Sauvageat, J.L., Raab, T. (2012). Hydrolysis of chicoric and caftaric acids with esterases and *Lactobacillus johnsonii* in vitro and in a gastrointestinal model. *J. Agric. Food Chem.* 60: 9236-9241.
- Bel-Rhliid, R., Crespy, V., Pagé-Zoerkler, N., Nagy, K., Raab, T. and Hansen, C-E. (2009). Hydrolysis of rosmarinic acid from Rosemary extract with esterases and *Lactobacillus johnsonii* in vitro and in a gastrointestinal model. *J. Agric. Food Chem.* 57: 7700-7705.
- Blanquet-Diot, S., Denis, S., Chalancon, S., Chaira, F., Cardot, J.-M., Alric, M. (2012). Use of artificial digestive systems to investigate the biopharmaceutical factors influencing the survival of probiotic yeast during gastrointestinal transit in humans. *Pharm. Research* 29: 1444-1453.
- Cordonnier, C., Thévenot, J., Etienne-Mesmin, L., Denis, S., Alric, M., Livrelli, V., Blanquet-Diot, S. (2015). Dynamic in vitro models of the human gastrointestinal tract as relevant tools to assess the survival of probiotic strains and their interactions with gut microbiota. *Microorganisms* 3: 725-745.
- Etienne-Mesmin, L., Livrelli, V., Privat, M., Denis, S., Cardot, J.M., Alric, M., Blanquet-Diot, S. (2011). Effect of a new probiotic *Saccharomyces cerevisiae* strain on survival of *Escherichia coli* O157:H7 in a dynamic gastrointestinal model. *Applied and Environmental Microbiology* 77: 1127-1131.
- Fernandez, B., Hammami, R., Savard, P., Jean, J., Fliss, I. (2013). *Pediococcus acidilactici* UL5 and *Lactococcus lactis* ATCC 11454 are able to survive and express their bacteriocin genes under simulated gastrointestinal conditions. *J. Appl. Microbiol.* 116: 677-688.
- Fujii, A., Crédoz, Y., Maathuis, A.J.H., Nishida, S. (2015). Different viability of probiotic strains accessible in Japanese dairy market demonstrated by *in vitro* methods. *Milk Sci.* 64 (2): 99-106.
- Gagnon, M., Savard, P., Rivièrè, A., LaPointe, G., Roy, D. (2014). Bioaccessible antioxidants in milk fermented by *Bifidobacterium longum* subsp. *longum* strains. *BioMed Res. Internat.*, ID 169381.

- Gänzle, M.G., Hertel, C., Van der Vossen, J.M.B.M. and Hammes, W.P. (1999). Effect of bacteriocin-producing lactobacilli on the survival of *Escherichia coli* and *Listeria* in a dynamic model of the stomach and the small intestine. *Int. J. Food Microbiology* 48: 21-35.
- Hanchi, H., Hammami, R., Kourda, R., Ben Hamida, J., Fliss, I. (2014). Bacteriocinogenic properties and in vitro probiotic potential of enterococci from Tunisian dairy products. *Arch. Microbiol.* 196: 331-344.
- Hatanaka, M., Nakamura, Y., Maathuis, A.J.H., Venema, K., Murota, I., Yamamoto, Y. (2012). Influence of *Bacillus subtilis* C-3102 on microbiota in a dynamic *in vitro* model of the gastrointestinal tract simulating human conditions. *Beneficial Microbes* 3 (3): 229-236.
- Havenaar, R. (1999). The model selection tool. *Dairy Industries International*, 64 (6): 33-36.
- Jedidi, H., Champagne, C.P., Raymond, Y., Farnworth, E., van Calsteren, M-R., Chouinard, P.Y., Fliss, I., (2014). Effect of milk enriched with conjugated linoleic acid and digested in a simulator (TIM-1) on the viability of probiotic bacteria. *Internat. Dairy J.* 37: 20-25.
- Keller, D., Van Dinter, R., Cash, H., Farmer, S., Venema, K. (2017). *Bacillus coagulans* GBI-30, 6086 increases plant protein digestion in a dynamic, computer-controlled *in vitro* model of the small intestine (TIM-1). *Beneficial Microbes* 8 (3): 491-496.
- Khalf, M., Dabour, N., Kheadr, E. and Fliss, I. (2010). Viability of probiotic bacteria in maple sap products under storage and gastrointestinal conditions. *Bioresource Technology* 101: 7966-7972.
- Kheadr, E., Zihler, A., Dabour, N., Lacroix, C., Le Blay, G. and Fliss, I. (2010). Study of the physicochemical and biological stability of pediocin PA-1 in the upper gastrointestinal tract conditions using a dynamic *in vitro* model. *J. Appl. Microbiol.* 109: 54-64.
- Kheadr, E.E., Dabour, N., Petit, G., Vuilleumard, J-C. (2011). Probiotic-delivering capacity of dairy products: *In vitro* assessment using a gastro-intestinal dynamic model. *Internat. J. Probiotics Prebiotics* 6 (2): 73-80.
- Maathuis, A., Keller, D. and Farmer, S. (2010). Survival and metabolic activity of the GanedenBC<sup>30</sup> strain of *Bacillus coagulans* in a dynamic *in vitro* model of the stomach and small intestine. *Beneficial Microbes* 1 (1): 31-36.
- Makivuokko, H., Wacklin, P., Koenen, ME., Laamanen, K., Alakulppi, N., Venema, K. and Matto, J. 2012. Isolation of bifidobacteria for blood group secretor status targeted personalised nutrition. *Microbial Ecology in Health & Disease* 23: 28-34.
- Marteau, P., Minekus, M., Havenaar, R. and Huis in 't Veld, J.H.J. (1997). Survival of lactic acid bacteria in a dynamic model of the stomach and small intestine: Validation and the effects of bile. *J. Dairy Sci.* 80: 1031-1037.
- Martinez, R.C.R., Aynaou, A-E., Albrecht, S., Schols, H.A., De Martinis, E.C.P., Zoetendal, E.G., Venema, K., Saad, S.M.I., Smidt, H. (2011). In vitro evaluation of gastrointestinal survival of *Lactobacillus amylovorus* DSM 16698 alone and in combination with galactooligosaccharides, milk and/or *Bifidobacterium animalis* subsp. *lactis* bb-12. *Internat. J. Food Microbiol.* 149: 152-158.
- Miettinen, M., Alander, M., von Wright, A., Vuopio-Varkila, J., Marteau, P., Huis in't Veld, J. and Mattila-Sandholm, T. (1998). The survival of and cytokine induction by lactic acid bacteria after passage through a gastrointestinal model. *Microbial Ecology Health Disease* 10: 141-147.
- Miszczucha, S.D., Thévenot, J., Denis, S., Callon, C., Livrelli, V., Alric, M., Montel, M-C., Blanquet-Diot, S., Thevenot-Sergentet, D. (2014). Survival of *Escherichia coli* O26:H11 exceeds that of *Escherichia coli* O157:H7 as assessed by simulated human digestion of contaminated raw milk cheeses. *Internat. J. Food Microbiol.* 172: 40-48.
- Roussel, C., Cordonnier, C., Galia, W., Le Goff, O., Thévenot, J., Chalancon, S., Alric, M., Thevenot-Sergentet, D., Leriche, F., Van de Wiele, T., Livrelli, V., Blanquet-Diot, S. (2016) Increased EHEC survival and virulence gene expression indicate an enhanced pathogenicity upon simulated pediatric gastrointestinal conditions. *Pediatric Res.* 80 (5): 734-743.
- Uriot, O., Galia, W., Ablavi Awussi, A., Perrin, C., Denis, S., Chalancon, S., Lorson, E., Poirson, C., Junjua, M., Le Roux, Y., Alric, M., Dary, A., Blanquet-Diot, S., Roussel, Y. (2016). Use of dynamic gastro-intestinal model TIM to explore the survival of the yogurt bacterium *Streptococcus thermophilus* and the metabolic activities induced in the simulated human gut. *Food Microbiology* 53: 18-29.

Zhu, Y., Havenaar, R. and Venema, K. (2011) Response to: Pitino et al. (2010). Food Microbiology 27:1121-1127 "Survival of *Lactobacillus rhamnosus* strains in the upper gastrointestinal tract". Food Microbiology, 28:1110.

## Micro-ecology and bioconversion studies in the large-intestinal

- Aguirre, M., Bussolo de Souza, C., Venema, K. (2016). The gut microbiota from lean and obese subjects contribute differently to the fermentation of arabinogalactan and inulin. PLOS one, July: 1-18.
- Aguirre, M., Eck, A., Koenen, M.E., Savelkoul, P.H.M., Budding, A.E., Venema, K. (2016). Diet drives quick changes in the metabolic activity and composition of human gut microbiota in a validated in vitro gut model. Res. Microbiol. 167: 114-125.
- Aguirre, M., Eck, A., Koenen, M.E., Savelkoul, P.H.M., Budding, A.E., Venema, K. (2015). Evaluation of an optimal preparation of human standardized fecal inocula for *in vitro* fermentation studies. J. Microbiol. Methods 117: 78-84.
- Aguirre, M., Jonkers, D.M.A.E., Troost, F., Roeselers, G., Venema, K. (2014), In vitro characterization of the impact of different substrates on metabolite production, energy extraction and composition of gut microbiota from lean and obese subjects. Plos One (Open Access) Nov. 26: 1-23 (DOI: 10.1371/journal.pone.0113864).
- Aguirre, M., Ramiro-Garcia, J., Koenen, M.E., Venema, K. (2014). To pool or not to pool? Impact of the use of individual and pooled fecal samples for *in vitro* fermentation studies. J. Microbiol. Methods 107: 1-7.
- Binsl, T.W., De Graaf, A.A., Venema, K., Heringa, J., Maathuis, M., De Waard, P., Van Beek, J.H.G.M. (2010). Measuring non-steady-state metabolic fluxes in starch-converting faecal microbiota *in vitro*. Beneficial Microbes, December 1 (4): 391-405.
- Bussolo de Souza, C., Roeselers, G., Troost, F., Jonkers, D., Koenen, M.E., Venema, K. (2014). Prebiotic effects of cassava bagasse in TNO's in vitro model of the colon in lean versus obese microbiota. J. Func. Food 11: 210-210. Corrigendum: J. Func. Foods 13 (2015): 391.
- De Graaf, A.A., Maathuis, A., de Waard, P., Deutz, N.E.P., Dijkema, C., de Vos, W.M. and Venema, K. (2010). Profiling human gut bacterial metabolism and its kinetics using [U-13C]glucose and NMR. NMR Biomed. 23: 2-12.
- De Graaf, A.A., Venema, K. (2008). Gaining insight into microbial physiology in the large intestine: A special Role for stable isotopes. Advances Microbial Physiol. 53: 73-170.
- Egert, M., de Graaf, A., Maathuis, A., de Waard, P., Plugge, C.M., Smidt, H., Deutz, N.E.P., Dijkema, C., de Vos, W.M., Venema, K. (2007). Identification of glucose-fermenting bacteria present in an in vitro model of the human intestine by RNA-stable isotope probing. FEMS Microb. Ecol. 60 (1): 126-135.
- Faessler, C, Arrigoni, E., Venema, K., Brouns, F. and Amado, R. (2006). *In vitro* fermentability of differently digested resistant starch preparations. Mol. Nutr. Food Res. 50: 1220-1228.
- Gao, K., Xu, A., Krul, C., Venema, K., Liu, Y., Niu, Y., Lu, J., Bensoussan, L., Seeram, N.P., Heber, D. and Henning, S.M. (2006). Of the major phenolic acids formed during human microbial fermentation of tea, citrus and soy flavonoids supplements, only 3,4-dihydroxyphenylacetic acid has antiproliferative activity. J. Nutr. 136: 52-57.
- Hatanaka, M., Nakamura, Y., Maathuis, A.J.H., Venema, K., Murota, I., Yamamoto, Y. (2012). Influence of *Bacillus subtilis* C-3102 on microbiota in a dynamic *in vitro* model of the gastrointestinal tract simulating human conditions. Beneficial Microbes 3 (3): 229-236. `
- Koenen, M.E., Cruz Rubio, J.M., Mueller, M., Venema, K. (2016). The effect of agave fructan products on the activity and composition of the microbiota determined in a dynamic *in vitro* model of the human proximal large intestine. J. Funct. Foods 22: 201-210.
- Kong, H., Wang, M., Venema, K., Maathuis, A., van der Heijden, R., van der Greef, J., Xu, G., Hankemeier, T. (2009). Bioconversion of red ginseng saponins in the gastro-intestinal tract *in vitro*

- model studied by high-performance liquid chromatography–high resolution Fourier transform ion cyclotron resonance mass spectrometry. *J. Chromatography A* 1216: 2195–2203.
- Kortman, G.A.M., Dutilh, B.E., Maathuis, A.J.H., Engelke, U.F., Boekhorst, J., Keegan, K.P., Nielsen, F.G.G., Betley, J., Weir, J.C., Kingsbury, Z., Kluijtmans, L.A.J., Swinkels, D.W., Venema, K., Tjalsma, H. (2016). Microbial metabolism shifts towards an adverse profile with supplementary iron in the TIM-2 *in vitro* model of the human colon. *Frontiers Microbiol.* 6, art. 1481. doi: 10.3389/fmicb.2015.01481.
- Kovatcheva-Datchary, P., Egert, M., Maathuis, A., Rajilic-Stojanovic, M., de Graaf, A., Smidt, H., de Vos, W. and Venema, K. (2009). Linking phylogenetic identities of bacteria to starch fermentation in an *in vitro* model of the large intestine by RNA-based stable isotope probing. *Environmental Microbiology*, 11 (4): 914–926.
- Krul, C., Humblot, C., Philippe, C., Vermeulen, M., Van Nuenen, M., Havenaar, R., Rabot, S. (2002). Metabolism of sinigrin (2-propenyl glucosinolate) by the human colonic microflora in a dynamic *in vitro* large-intestinal model. *Carcinogenesis* 23 (6): 1009-1016.
- Lamers, R.-J., Wessels, E., Van de Sandt, J., Venema, K., Schaafsma, G., Van der Greef, J. and Van Nesselrooij J. (2003). A pilot study to investigate effects of inulin on Caco-2 cells through *in vitro* metabolic fingerprinting. *J. Nutr.* 133: 3080-3084.
- Maathuis, A.J.H., van den Heuvel, E.G., Schoterman, M.H.C., Venema, K. 2012. Galacto-Oligosaccharides have prebiotic activity in a dynamic *in vitro* colon model using a <sup>13</sup>C-labeling technique. *J. Nutrition* 142 (7): 1205-1212.
- Maathuis, A., Hoffman, A., Evans, A., Sanders, L., Venema, K. (2009). The Effect of the undigested fraction of maize products on the activity and composition of the microbiota determined in a dynamic *in vitro* model of the human proximal large intestine. *J. American College of Nutrition*, Vol. 28 (6): 657–666.
- Martinez, R.C.R., Cardarelli, H.R., Borst, W., Albrecht, S., Schols, H., Gutierrez, O.P., Maathuis, A.J.H., de Melo Franco, B.D.G., De Martinis, E.C.P., Zoetendal, E.G., Venema, K., Saad, S.M.I., Smidt, H. (2013). Effect of galactooligosaccharides and *Bifidobacterium animalis* Bb-12 on growth of *Lactobacillus amylovorus* DSM 16698, microbial community structure, and metabolic production in an *in vitro* colonic model set up with human or pig microbiota. *FEMS Microbiol. Ecol.* 84: 110-123.
- Motelica-Wagenaar, A.M., Nauta, A., van den Heuvel, E.G.H.M., Kleerebezem, R. (2014). Flux analysis of the human proximal colon using anaerobic digestion model 1. *Anaerobe* 28: 137-148.
- Rajilic-Stojanovic, M., Maathuis, A., Heilig, H., Venema, K., De Vos, W., Smidt, H. (2010). Evaluating the microbial diversity of an *in vitro* model of the human large intestine by phylogenetic microarray analysis. *Microbiology* 156: 3270-3281.
- Ramasamy, U.S., Venema, K., Schols, H.A., Gruppen, H. (2014). Effect of soluble and insoluble fibers within the *in vitro* fermentation of chicory root pulp by human gut bacteria. *J. Agr. Food Chem.* 62: 6794-6802.
- Ramasamy, U.S., Venema, K., Gruppen, H., Schols, H.A. (2014). The fate of chicory root pulp polysaccharides during fermentation in the TNO *in vitro* model of the colon (TIM-2). *Bioact. Carb. Dietary Fibre* 4: 48-57.
- Rehman, A., Heinsen, F.-A., Koenen, M.E., Venema, K., Knecht, H., Hellmig, S., Schreiber, S., Ott, S.J. 2012. Effects of probiotics and antibiotics on the intestinal homeostasis in a computer controlled model of the large intestine. *MBC Microbiology* 12, 47: 1-10.
- Reimer, R.A., Maathuis, A.J.H., Venema, K., Lyon, M.R., Gahler, R.J., Wood, S. (2014). Effect of the novel polysaccharide PolyGlycopleX® on short-chain fatty acid production in a computer-controlled *in vitro* model of the human large intestine. *Nutrients* 6: 1115-1127. (doi: 10-3390/nu6031115).
- Rose, D.J., Venema, K., Keshavarzian, A. and Hamaker, B.R. (2010). Starch-entrapped microspheres show a beneficial fermentation profile and decrease in potentially harmful bacteria during *in vitro* fermentation in faecal microbiota obtained from patients with inflammatory bowel disease. *British J. Nutr.* 103: 1514-1524.

- Tabernero, M., Venema, K., Maathuis, A.J.H., Saura-Calixto, F.D. (2011). Metabolite production during *in vitro* colonic fermentation of dietary fiber: Analysis and comparison of two European diets. *J. Agricult. Food Chemistry* 59 (16): 8968-8975.
- Van den Abbeele, P., Venema, K., Van de Wiele, T., Verstraete, W., Possemiers, S. (2013). Different human gut models reveal the distinct fermentation patterns of arabinoxylan versus inulin. *J. Agric. Food Chem.* 61: 9819-9827.
- Van der Vossen, J.M.B.M., Havekes, W.A.L.M., Koster, D.S., Ten Brink, B., Minekus, M., Havenaar, R., Huis in 't Veld, J.H.J., Overeem, J., Hendriks, N., Hofstra, H. (1998). Development and application of *in vitro* intestinal tract model for safety evaluation of genetically modified foods. In: *Food safety Evaluation of genetically modified foods as a basis for market introduction*. Ministry of Economic Affairs, The Hague, The Netherlands
- Van der Werf, M.J. and Venema, K. (2001). Bifidobacteria: genetic modification and the study of their role in the colon. *J. Agric. Food Chem.* 49: 378-383.
- Van Nuenen, M., de Ligt, R.A., Doornbos, R.P., Venema, K., Van der Woude, C.J., and Kuipers, E.J. (2005). The influence of microbial metabolites on human intestinal epithelial cells and macrophages *in vitro*. *FEMS Immunol Med Microbiol* 45 (2): 183-189.
- Van Nuenen, H.M.C., Venema, K., Van der Woude, J.C.J. and Kuipers, E.J. (2004). The metabolic activity of fecal microbiota from healthy individuals and patients with inflammatory bowel disease. *Digestive Disease and Sciences* 49 (3): 485-491.
- Van Nuenen, H.M.C., Meyer, P.D., Venema, K. (2003). The effect of various inulins and *Clostridium difficile* on the metabolic activity of the human colonic microbiota *in vitro*. *Microbial Ecology in Health and Disease* 15 (2-3): 137-144.
- Venema K. 2015. Health Effects of Pro and Prebiotics: Utilization of Sophisticated *In Vitro* Tools. In: *Beneficial Microorganisms in Medical and Health Applications; Series: Microbiology Monographs, Vol. 28; M.-T. Liong (Ed.) 2015. ISBN: 978-3-319-23212-6. Chapter 1; pp. 1-18.*
- Venema, K. (2014). *In vitro* assessment of the bioactivity of food oligosaccharides. In: *Food Oligosaccharides: Production, Analysis and Bioactivity*. Eds.: F.J. Moreno & M. Luz Sanz. John Wiley & Sons, Ltd.
- Venema, K., Van den Abbeele, P. (2013). Experimental models of the gut microbiome. *Best Practice & Research Clinical Gastroenterology* 27: 115-126.
- Venema, K., Vermunt, S.H.F. and Brink, E.J. (2005). D-Tagatose increases butyrate production by colonic microbiota in healthy men and women. *Microbial Ecology Health Dis.* 17: 47-57.
- Venema, K. and Maathuis, A. (2003). A PCR-based method for identification of bifidobacteria from the human alimentary tract at the species level. *FEMS Microbiol. Letters* 224 (1): 143-149.
- Venema, K. and Van de Sandt, H. (2003). Interaction between food components, intestinal microbiota and intestinal mucosa as a function of intestinal health. *AgroFoods* 14 (March/April): 62-66.
- Venema, K., Van Nesselrooij, J., Lamers, R.-J. and Van de Sandt, J. (2003). Metabolic fingerprinting of Caco-2 cells. Effect of inulin and its fermentative metabolites. *Nutrafoods* 2 (1): 5-12.
- Venema, K., Van Nuenen, H.M.C., Van den Heuvel, E.G., Pool, W., Van der Vossen, J.M.B.M. (2003). The effect of lactulose on the composition of the intestinal microbiota and short-chain fatty acid production in human volunteers and a computer-controlled model of the proximal large intestine. *Microbial Ecology in Health and Disease*, 15 (2-3): 94-105.
- Venema, K., Van Nuenen, H.M.C., Smeets-Peeters, M.J.E., Minekus, M. and Havenaar, R. (2000). TNO's *in vitro* large intestinal model: an excellent screening tool for functional food and pharmaceutical research. *Ernährung/Nutrition* 24 (12): 558-564.

## (Food) Safety - Chemicals / Toxicology / Toxin binding / Polluted soil

- Avantaggiato, G., Havenaar, R. and Visconti, A. (2007). Assessment of the multi-mycotoxin binding efficacy of a carbon/aluminosilicate based product in an *in vitro* gastrointestinal model. *J. Agricultural Food Chem.* 55: 4810-4819.

- Avantaggiato, G., Havenaar, R. and Visconti, A. (2004). Evaluation of the intestinal absorption of deoxynivalenol and nivalenol by an *in vitro* gastrointestinal model, and the binding efficacy of activated carbon and other adsorbent materials. *Food Chemical Tox.* 42 (5): 817-824.
- Avantaggiato, G., Havenaar, R. and Visconti, A. (2003). Assessing the zearalenone binding activity of adsorbent materials during passage through a dynamic gastrointestinal model. *Food Chemical Tox.* 41: 1283-1290.
- Blanquet S, Meunier, JP, Minekus M, Marol-Bonnin S and Alric M (2003). Recombinant *Saccharomyces cerevisiae* expressing a P450 in artificial digestive systems: a model for biotransformation in the human digestive environment. *Appl. Env. Microbiol.* 69: 2884-2892.
- Bockting, G. Van der Valk, W. (1998). Gastrointestinale absorptie van lood uit verontreinigde grond. *Bodem 2*: 74-76.
- Dominy, N.J., Davoust, E. and Minekus, M. (2004). Adaptive function of soil consumption: an *in vitro* study modeling the human stomach and small intestine. *J. Exp. Biol.* 207: 319-324.
- Frank, N., Dubois, M., Scholz, G., Seefelder, W., Chuat, J-Y., Schilter, B. (2013). Application of gastrointestinal modelling to the study of the digestion and transformation of dietary glycidyl esters. *Food Add. Containm. Part A*, 30 (1): 69-79. Doi: 10.1080/19440049.2012.732245.
- Havenaar, R., de Jong, A., Koenen, M.J., van Bilsen, J., Janssen, A.M., Labij, E., Westerbeek, H.J.M. (2013). Digestibility of transglutaminase cross-linked caseinate versus native caseinate in an *in vitro* multi-compartmental model simulating young child and adult gastrointestinal conditions. *J. Agric. Food Chem.* (accepted).
- Helou, C., Denis, S., Spatz, M., Marier, D., Rame, V., Alric, M., Tessier, F.J., Gadonna-Widehem, P. (2015). Insights into bread melanoidins: Fate in the upper digestive tract and impact on the gut microbiota using *in vitro* systems. *Food Func.* 6 (12): 3737-3745.
- Krul, C.A.M., Zeilmaier, M., Schothorst, R. and Havenaar, R. (2004). Intra-gastric formation and modulation of N-nitrosodimethylamine in a dynamic *in vitro* gastrointestinal model under human physiological conditions. *Food Chem. Toxicology* 42: 51-63.
- Krul, C.A.M., Luiten-Schuite, A., Baan, R., Verhagen, H., Mohn, G., Feron, V., and Havenaar, R. (2000). Application of a dynamic *in vitro* gastrointestinal tract model to study the availability of food mutagens, using heterocyclic aromatic amines as model compounds. *Food and Chemical Toxicology* (38): 783-792.
- Lefebvre, D.E., Venema, K., Gombau, L., Valerio Jr, L.G., Raju, J., Genevieve S. Bondy, G.S., Bouwmeester, H., Singh, R.P., Clippinger, A.J., Collnot, E.-M., Mehta, R., Stone, V. (2014). Utility of models of the gastrointestinal tract for assessment of the digestion and absorption of engineered nanomaterials released from food matrices. *Nanotoxicology* (early on line 1-20; DOI: 10.3109/17435390.2014.948091).
- Oomen, A.G.; Hack, A.; Minekus, M.; Zeijdner, E.; Cornelis, C.; Schoeters, G.; Verstraete, W.; Wiele, T. van de; Wragg, J.; Rempelberg, C.J.M.; Sips, A.J.A.M.; Wijnen, J.H. van. (2002). Comparison of five *in vitro* digestion models to study the bioaccessibility of soil contaminants. *Environmental Science & Technology* 36 (15): 3326-3334.
- Roussel, C., Cordonnier, C., Galia, W., Le Goff, O., Thévenot, J., Chalancon, S., Alric, M., Thevenot-Sergentet, D., Leriche, F., Van de Wiele, T., Livrelli, V., Blanquet-Diot, S. (2016) Increased EHEC survival and virulence gene expression indicate an enhanced pathogenicity upon simulated pediatric gastrointestinal conditions. *Pediatric Res.* 80 (5): 734-743.
- Torres-Escribano, S., Denis, S., Blanquet-Diot, S., Galatayud, M., Barrios, L., Velez, D, Alric, M., and Montoro, R. (2011). Comparison of a static and a dynamic *in vitro* model to estimate the bioaccessibility of As, Cd, Pb and Hg from food reference materials *Fucus* sp. (IAEA-140/TM) and lobster hepatopancreas (TORT-2). *Sci. Total Environment* 409: 604-611.
- Van de Wiele, T., Oomen, A., Wragg, J., Cave, M., Minekus, M., Hack, A., Cornelis, C., Rempelberg, C., De Zwart, L., Klinck, B., Van Wijnen, J., Verstraete, W., and Sips, A. (2007). Comparison of five *in vitro* digestion models to *in vivo* experimental results: Lead bioaccessibility in the human gastrointestinal tract. *J. Experimental Sci. Health. Part A*, 42: 1203-1211.



Zeilmaker, M.J., Bakker, M.I., Schothorst, R.; Slob, W. (2010). Risk assessment of N-nitrosodimethylamine formed endogenously after fish-with-vegetable meals. *Toxicological Sciences* 116 (1): 323-335.